

AD-A071 694

MICHIGAN UNIV ANN ARBOR SCHOOL OF MEDICINE

F/G 6/5

POPULATION GENETICS OF HEMOGLOBIN E, THALASSEMIA AND RELATED GE--ETC(U)

MAR 78 D L RUCKNAGEL

DA-49-193-MD-2847

NL

UNCLASSIFIED


| OF |

AD  
A071694



END  
DATE  
FILMED  
8-79

DDC



AD A 071 694

DDC FILE COPY

LEVEL

10/17/78

1

SC

Population Genetics of Hemoglobin E, Thalassemia  
and Related Genetic Polymorphisms in Thailand

Final Report  
(1965 - 1970)

March 1978

by

Donald L. Rucknagel, M. D.

Supported by

US Army Medical Research and Development Command  
Fort Detrick, Frederick, Maryland 21701

Contract No. DA-49-193-MD-2847

University of Michigan Medical School  
Ann Arbor, Michigan 48104

X School of medicine

DDC

JUL 26 1979

Approved for public release; distribution unlimited

The findings in this report are not to be construed  
as an official Department of the Army position unless  
so designated by other authorized documents.

228 850

79 07 17 080

LB

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9)
4. TITLE (and Subtitle) Population Genetics of Hemoglobin E, Thalassemia and Related Genetic Polymorphisms in Thailand.		5. TYPE OF REPORT & PERIOD COVERED Final Report 1965 - 1970
6. AUTHOR(s) (10) Donald L. Rucknagel, M. D.		7. PERFORMING ORG. REPORT NUMBER
8. PERFORMING ORGANIZATION NAME AND ADDRESS University of Michigan Medical School Ann Arbor, Michigan 48104		9. CONTRACT OR GRANT NUMBER(s) (15) DA-49-193-MD-2847
10. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick, Frederick, Maryland 21701		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (16) 61101A 3A061101A91C 00, 123
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 8p.		13. REPORT DATE (11) March 1978
		14. NUMBER OF PAGES 8 pages
		15. SECURITY CLASS. (of this report) Unclassified
		16. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
18. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
19. SUPPLEMENTARY NOTES		
20. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
21. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

The Thai population data have been analyzed using the following general approach. The phenotype frequency was determined for all of the genes after separating the Thais from hill tribesmen. From the total Thai sample we then removed first the children and then related adults. The frequency of genetic traits in the total or corrected samples did not differ significantly. These manipulations were necessary to deal with the randomness of the samples, because even truly random samples of small endogamous populations will contain relatives. Since we did not know precisely the mating structure of our populations we had no other more direct approach to the question of whether we had ascertained an inordinate number of relatives. We then calculated gene frequencies for each genetic system in each village and in each province and compared them statistically and for adherence to the Hardy-Weinberg equilibrium.

The anthropometrics and other quantitative variables such as uric acid and serum iron levels were analyzed by regressing onto age and then comparing between groups by covariance analysis. Group comparisons included comparisons between provinces in Thailand, between sexes, and for attributes such as serum iron and hemoglobin levels between the hemoglobin E, thalassemia, and normal phenotypes.

In view of the complexity of the study a table is appended to demonstrate some of the findings. First, there is a great deal of heterogeneity in gene frequency within provinces in Thailand (not shown in the table) which cannot be attributed to small sample size. Despite this variation, gene frequencies appear to vary systematically in the country as judged from the ten provinces which we have studied. In Northeastern Thailand, in addition to the high frequency of Hb E, the Thai electrophoretic variant on ceruloplasmin, transferrin D, blood groups Ms, and secretor are more frequent than in other areas; Lewis (a+b-), and 6-phosphogluconate dehydrogenase type B are somewhat less frequent. Frequencies in Surin are strikingly out of line with those elsewhere in Northeastern Thailand, suggesting that Cambodians are somewhat different. For instance, the frequencies of ceruloplasmin Thai, secretor, Lewis (a-b-), and blood group B are greater; blood groups O and Ns are less frequent than elsewhere in this region.

Accession For	NTIS GRA&I	<input checked="" type="checkbox"/>	
DDC TAB	Unannounced		
Justification			
By			
Distribution			
Availability Codes			
Avail and/or			
Dist special			
Dist			A



In Northern Thailand the beta-thalassemia gene is most frequent. This does not appear to be derived from the hill tribesmen inasmuch as it was nearly absent from our small series of tribesmen. This region is characterized by elevated levels of glucose-6-phosphate dehydrogenase deficiency, Diego blood group, and Ms, the lower frequencies of 6-phosphogluconate type B, secretor, blood group B, and MS.

Southern Thailand is characterized by an elevated frequency of MS, and lower frequencies of hemoglobin E, G-6-PD deficiency, Ms, and Lewis (a-b-) than in most other areas.

The analysis of anthropometric measurements shows a large number of non-systematic differences between provinces.

Some systematic differences are apparent, however. For instance, individuals in Northern Thailand appear significantly smaller than those examined elsewhere in the country. Interpretation of these differences will be difficult, until statistical inter-racial distance measurements are completed; these are in progress.

The differences in gene frequencies outlined suggest that genetic drift and founder effect play a large role in so-called microdifferentiation within provinces. Whereas the differences in gene frequencies in the various provinces do not correlate perfectly, the fact that so many gene frequencies vary so much from one region to another suggests that this variation is more likely to be due to gene flow or migration rather than differential selective values throughout the country. If, for instance, the high frequency of thalassemia were the only characteristic of the Northern Thai, malaria selection pressure would be a reasonable explanation. If a rational relationship were apparent for several other genetic characteristics in this area, one might still invoke selection. For instance, malaria might select for genetic mutants of the hemoglobin, glucose-6-phosphate dehydrogenase, and haptoglobin loci since all three of these proteins are involved with the integrity of the red blood cell. We have no rational basis for invoking entities such as the blood groups, for instance, also in a selection model. Migration seems a more likely common denominator.

4 The gene frequency data have been collated, and the terminal report is being prepared on the descriptive aspects of the study. The fertility data have been analyzed in a preliminary fashion but set aside for the moment because they present special problems in analysis for which we have not as yet worked out the methodology. 7

	B	O	M <sub>s</sub>	Df. <sup>a</sup>	Sec.	Lc (e+b)
HbE	β-Th-1	Gc-1	Hd-1	Cp-Th		
	Tf-D					
6-PGD						

[illegible]



**Project 3A061101A91C IN-HOUSE LABORATORY INDEPENDENT RESEARCH**

**Task 00, In-House Laboratory Independent Research**

**Work Unit 123, Population genetics of hemoglobin E,  
thalassemia, and related genetic poly-  
morphisms in Thailand**

**Literature Cited.**

**1. Reference**

Shokeir, M. A., Rucknagel, D. L., Na-Nakorn, S., Wasi, P.,  
and Shreffler, D. C.: Genetic polymorphism of human  
serum ceruloplasmin in Thailand. Amer. J. Hum. Genet.,  
in press.

**2. Publications**

**None**

**12 Copies**

**Director (ATTN: SGRD-UWZ-AG)  
Walter Reed Army Institute of  
Research  
Walter Reed Army Medical Center  
Washington, D. C. 20012**

**4 Copies**

**HQDA (SGRD-AJ)  
Fort Detrick  
Frederick, MD 21701**

**12 Copies**

**Defense Documentation Center  
ATTN: DDC-TCA  
Cameron Station  
Alexandria, Virginia 22314**

**1 Copy**

**Dean  
School of Medicine  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, Maryland 20014**

**1 Copy**

**Superintendent  
Academy of Health Sciences, US Army  
ATTN: AHS-COM  
Fort Sam Houston, Texas 78234**